


COVID-19 Vaccine Timing by Age

Routine Schedule



^ An [8-week interval](#) may be preferable for some people, especially for males 12-39 years.






* Although use of mRNA COVID-19 vaccines is preferred, the Janssen vaccine may be offered in [some situations](#).

** People who received J&J for their primary and first booster dose may consider receiving an mRNA vaccine as a second booster.

View [Interim Clinical Considerations for Use of COVID-19 Vaccines](#) for details. Schedule is subject to change.

COVID-19 Vaccine Timing by Age

Schedule if Moderately or Severely Immunocompromised

Age	Vaccine	Primary Doses	1st Booster Dose	2nd Booster Dose	
5-11	Pfizer–Pediatric (5-11)		Booster		
12+	Pfizer/Comirnaty (12+)		1st Booster Ages 12-17: Pfizer 18+: Pfizer/Moderna (mRNA preferred) or J&J*	2nd Booster Ages 12-17: Pfizer 18+: Moderna/Pfizer	
18+	Moderna/Spikevax				
18+	Johnson & Johnson Pfizer/Moderna preferred*				

*Although use of mRNA COVID-19 vaccines is preferred, the Janssen vaccine may be offered in [some situations](#).

View [COVID-19 Vaccines for Moderately or Severely Immunocompromised People](#) for details. Schedule is subject to change.